

Inversion of the Dual Dunkl–Sonine Transform on $\mathbb{R}$ Using Dunkl Wavelets

Mohamed Ali MOUROU

Department of Mathematics, Faculty of Sciences, Al-Jouf University,
P.O. Box 2014, Al-Jouf, Skaka, Saudi Arabia
E-mail: mohamed_ali.mourou@yahoo.fr

Received March 02, 2009, in final form July 04, 2009; Published online July 14, 2009

doi:10.3842/SIGMA.2009.071

Abstract. We prove a Calderón reproducing formula for the Dunkl continuous wavelet transform on $\mathbb{R}$. We apply this result to derive new inversion formulas for the dual Dunkl–Sonine integral transform.

Key words: Dunkl continuous wavelet transform; Calderón reproducing formula; dual Dunkl–Sonine integral transform

2000 Mathematics Subject Classification: 42B20; 42C15; 44A15; 44A35

1 Introduction

The one-dimensional Dunkl kernel e_γ , $\gamma > -1/2$, is defined by

$$e_\gamma(z) = j_\gamma(iz) + \frac{z}{2(\gamma+1)} j_{\gamma+1}(iz), \quad z \in \mathbb{C},$$

where

$$j_\gamma(z) = \Gamma(\gamma+1) \sum_{n=0}^{\infty} \frac{(-1)^n (z/2)^{2n}}{n! \Gamma(n+\gamma+1)}$$

is the normalized spherical Bessel function of index γ . It is well-known (see [3]) that the functions $e_\gamma(\lambda \cdot)$, $\lambda \in \mathbb{C}$, are solutions of the differential-difference equation

$$\Lambda_\gamma u = \lambda u, \quad u(0) = 1,$$

where

$$\Lambda_\gamma f(x) = f'(x) + \left(\gamma + \frac{1}{2}\right) \frac{f(x) - f(-x)}{x}$$

is the Dunkl operator with parameter $\gamma+1/2$ associated with the reflection group $\mathbb{Z}_2$ on $\mathbb{R}$. Those operators were introduced and studied by Dunkl [2, 3, 4] in connection with a generalization of the classical theory of spherical harmonics. Besides its mathematical interest, the Dunkl operator Λ_α has quantum-mechanical applications; it is naturally involved in the study of one-dimensional harmonic oscillators governed by Wigner's commutation rules [6, 11, 16].

It is known, see for example [14, 15], that the Dunkl kernels on $\mathbb{R}$ possess the following Sonine type integral representation

$$e_\beta(\lambda x) = \int_{-|x|}^{|x|} \mathcal{K}_{\alpha,\beta}(x, y) e_\alpha(\lambda y) |y|^{2\alpha+1} dy, \quad \lambda \in \mathbb{C}, \quad x \neq 0, \quad (1.1)$$

where

$$\mathcal{K}_{\alpha,\beta}(x,y) := \begin{cases} a_{\alpha,\beta} \operatorname{sgn}(x) (x+y) \frac{(x^2 - y^2)^{\beta-\alpha-1}}{|x|^{2\beta+1}} & \text{if } |y| < |x|, \\ 0 & \text{if } |y| \geq |x|, \end{cases} \quad (1.2)$$

with $\beta > \alpha > -1/2$, and

$$a_{\alpha,\beta} := \frac{\Gamma(\beta+1)}{\Gamma(\alpha+1)\Gamma(\beta-\alpha)}.$$

Define the Dunkl–Sonine integral transform $\mathcal{X}_{\alpha,\beta}$ and its dual ${}^t\mathcal{X}_{\alpha,\beta}$, respectively, by

$$\begin{aligned} \mathcal{X}_{\alpha,\beta} f(x) &= \int_{-|x|}^{|x|} \mathcal{K}_{\alpha,\beta}(x,y) f(y) |y|^{2\alpha+1} dy, \\ {}^t\mathcal{X}_{\alpha,\beta} f(y) &= \int_{|x| \geq |y|} \mathcal{K}_{\alpha,\beta}(x,y) f(x) |x|^{2\beta+1} dx. \end{aligned}$$

Soltani has showed in [14] that the dual Dunkl–Sonine integral transform ${}^t\mathcal{X}_{\alpha,\beta}$ is a transmutation operator between Λ_α and Λ_β on the Schwartz space $\mathcal{S}(\mathbb{R})$, i.e., it is an automorphism of $\mathcal{S}(\mathbb{R})$ satisfying the intertwining relation

$${}^t\mathcal{X}_{\alpha,\beta} \Lambda_\beta f = \Lambda_\alpha {}^t\mathcal{X}_{\alpha,\beta} f, \quad f \in \mathcal{S}(\mathbb{R}).$$

The same author [14] has obtained inversion formulas for the transform ${}^t\mathcal{X}_{\alpha,\beta}$ involving pseudo-differential-difference operators and only valid on a restricted subspace of $\mathcal{S}(\mathbb{R})$.

The purpose of this paper is to investigate the use of Dunkl wavelets (see [5]) to derive a new inversion of the dual Dunkl–Sonine transform on some Lebesgue spaces. For other applications of wavelet type transforms to inverse problems we refer the reader to [7, 8] and the references therein.

The content of this article is as follows. In Section 2 we recall some basic harmonic analysis results related to the Dunkl operator. In Section 3 we list some basic properties of the Dunkl–Sonine integral transform and its dual. In Section 4 we give the definition of the Dunkl continuous wavelet transform and we establish for this transform a Calderón formula. By combining the results of the two previous sections, we obtain in Section 5 two new inversion formulas for the dual Dunkl–Sonine integral transform.

2 Preliminaries

Note 2.1. Throughout this section assume $\gamma > -1/2$. Define $L^p(\mathbb{R}, |x|^{2\gamma+1} dx)$, $1 \leq p \leq \infty$, as the class of measurable functions f on $\mathbb{R}$ for which $\|f\|_{p,\gamma} < \infty$, where

$$\|f\|_{p,\gamma} = \left(\int_{\mathbb{R}} |f(x)|^p |x|^{2\gamma+1} dx \right)^{1/p}, \quad \text{if } p < \infty,$$

and $\|f\|_{\infty,\gamma} = \|f\|_{\infty} = \operatorname{ess\,sup}_{x \in \mathbb{R}} |f(x)|$. $\mathcal{S}(\mathbb{R})$ stands for the usual Schwartz space.

The Dunkl transform of order $\gamma + 1/2$ on $\mathbb{R}$ is defined for a function f in $L^1(\mathbb{R}, |x|^{2\gamma+1} dx)$ by

$$\mathcal{F}_\gamma f(\lambda) = \int_{\mathbb{R}} f(x) e_\gamma(-i\lambda x) |x|^{2\gamma+1} dx, \quad \lambda \in \mathbb{R}. \quad (2.1)$$

Remark 2.2. It is known that the Dunkl transform $\mathcal{F}_\gamma$ maps continuously and injectively $L^1(\mathbb{R}, |x|^{2\gamma+1}dx)$ into the space $\mathcal{C}_0(\mathbb{R})$ (of continuous functions on $\mathbb{R}$ vanishing at infinity).

Two standard results about the Dunkl transform $\mathcal{F}_\gamma$ are as follows.

Theorem 2.3 (see [1]).

(i) For every $f \in L^1 \cap L^2(\mathbb{R}, |x|^{2\gamma+1}dx)$ we have the Plancherel formula

$$\int_{\mathbb{R}} |f(x)|^2 |x|^{2\gamma+1} dx = m_\gamma \int_{\mathbb{R}} |\mathcal{F}_\gamma f(\lambda)|^2 |\lambda|^{2\gamma+1} d\lambda,$$

where

$$m_\gamma = \frac{1}{2^{2\gamma+2}(\Gamma(\gamma+1))^2}. \quad (2.2)$$

(ii) The Dunkl transform $\mathcal{F}_\alpha$ extends uniquely to an isometric isomorphism from $L^2(\mathbb{R}, |x|^{2\gamma+1}dx)$ onto $L^2(\mathbb{R}, m_\gamma |\lambda|^{2\gamma+1}d\lambda)$. The inverse transform is given by

$$\mathcal{F}_\gamma^{-1}g(x) = m_\gamma \int_{\mathbb{R}} g(\lambda) e_\gamma(i\lambda x) |\lambda|^{2\gamma+1} d\lambda,$$

where the integral converges in $L^2(\mathbb{R}, |x|^{2\gamma+1}dx)$.

Theorem 2.4 (see [1]). The Dunkl transform $\mathcal{F}_\alpha$ is an automorphism of $\mathcal{S}(\mathbb{R})$.

An outstanding result about Dunkl kernels on $\mathbb{R}$ (see [12]) is the product formula

$$e_\gamma(\lambda x) e_\gamma(\lambda y) = T_\gamma^x(e_\gamma(\lambda \cdot))(y), \quad \lambda \in \mathbb{C}, \quad x, y \in \mathbb{R},$$

where T_γ^x stand for the Dunkl translation operators defined by

$$\begin{aligned} T_\gamma^x f(y) &= \frac{1}{2} \int_{-1}^1 f\left(\sqrt{x^2 + y^2 - 2xyt}\right) \left(1 + \frac{x-y}{\sqrt{x^2 + y^2 - 2xyt}}\right) W_\gamma(t) dt \\ &\quad + \frac{1}{2} \int_{-1}^1 f\left(-\sqrt{x^2 + y^2 - 2xyt}\right) \left(1 - \frac{x-y}{\sqrt{x^2 + y^2 - 2xyt}}\right) W_\gamma(t) dt, \end{aligned} \quad (2.3)$$

with

$$W_\gamma(t) = \frac{\Gamma(\gamma+1)}{\sqrt{\pi} \Gamma(\gamma+1/2)} (1+t) (1-t^2)^{\gamma-1/2}.$$

The Dunkl convolution of two functions f, g on $\mathbb{R}$ is defined by the relation

$$f *_\gamma g(x) = \int_{\mathbb{R}} T_\gamma^x f(-y) g(y) |y|^{2\gamma+1} dy. \quad (2.4)$$

Proposition 2.5 (see [13]).

(i) Let $p, q, r \in [1, \infty]$ such that $\frac{1}{p} + \frac{1}{q} - 1 = \frac{1}{r}$. If $f \in L^p(\mathbb{R}, |x|^{2\gamma+1}dx)$ and $g \in L^q(\mathbb{R}, |x|^{2\gamma+1}dx)$, then $f *_\gamma g \in L^r(\mathbb{R}, |x|^{2\gamma+1}dx)$ and

$$\|f *_\gamma g\|_{r,\gamma} \leq 4 \|f\|_{p,\gamma} \|g\|_{q,\gamma}. \quad (2.5)$$

(ii) For $f \in L^1(\mathbb{R}, |x|^{2\gamma+1}dx)$ and $g \in L^p(\mathbb{R}, |x|^{2\gamma+1}dx)$, $p = 1$ or 2 , we have

$$\mathcal{F}_\gamma(f *_\gamma g) = \mathcal{F}_\gamma f \mathcal{F}_\gamma g. \quad (2.6)$$

For more details about harmonic analysis related to the Dunkl operator on $\mathbb{R}$ the reader is referred, for example, to [9, 10].

3 The dual Dunkl–Sonine integral transform

Throughout this section assume $\beta > \alpha > -1/2$.

Definition 3.1 (see [14]). The dual Dunkl–Sonine integral transform ${}^t\mathcal{X}_{\alpha,\beta}$ is defined for smooth functions on $\mathbb{R}$ by

$${}^t\mathcal{X}_{\alpha,\beta}f(y) := \int_{|x| \geq |y|} \mathcal{K}_{\alpha,\beta}(x, y) f(x) |x|^{2\beta+1} dx, \quad y \in \mathbb{R}, \quad (3.1)$$

where $\mathcal{K}_{\alpha,\beta}$ is the kernel given by (1.2).

Remark 3.2. Clearly, if $\text{supp}(f) \subset [-a, a]$ then $\text{supp}({}^t\mathcal{X}_{\alpha,\beta}f) \subset [-a, a]$.

The next statement provides formulas relating harmonic analysis tools tied to Λ_α with those tied to Λ_β , and involving the operator ${}^t\mathcal{X}_{\alpha,\beta}$.

Proposition 3.3.

- (i) The dual Dunkl–Sonine integral transform ${}^t\mathcal{X}_{\alpha,\beta}$ maps $L^1(\mathbb{R}, |x|^{2\beta+1} dx)$ continuously into $L^1(\mathbb{R}, |x|^{2\alpha+1} dx)$.
- (ii) For every $f \in L^1(\mathbb{R}, |x|^{2\beta+1} dx)$ we have the identity

$$\mathcal{F}_\beta(f) = \mathcal{F}_\alpha \circ {}^t\mathcal{X}_{\alpha,\beta}(f). \quad (3.2)$$

- (iii) Let $f, g \in L^1(\mathbb{R}, |x|^{2\beta+1} dx)$. Then

$${}^t\mathcal{X}_{\alpha,\beta}(f *_\beta g) = {}^t\mathcal{X}_{\alpha,\beta}f *_\alpha {}^t\mathcal{X}_{\alpha,\beta}g. \quad (3.3)$$

Proof. Let $f \in L^1(\mathbb{R}, |x|^{2\beta+1} dx)$. By Fubini's theorem we have

$$\begin{aligned} \int_{\mathbb{R}} {}^t\mathcal{X}_{\alpha,\beta}(|f|)(y) |y|^{2\alpha+1} dy &= \int_{\mathbb{R}} \left(\int_{|x| \geq |y|} \mathcal{K}_{\alpha,\beta}(x, y) |f(x)| |x|^{2\beta+1} dx \right) |y|^{2\alpha+1} dy \\ &= \int_{\mathbb{R}} |f(x)| \left(\int_{-|x|}^{|x|} \mathcal{K}_{\alpha,\beta}(x, y) |y|^{2\alpha+1} dy \right) |x|^{2\beta+1} dx. \end{aligned}$$

But by (1.1),

$$\int_{-|x|}^{|x|} \mathcal{K}_{\alpha,\beta}(x, y) |y|^{2\alpha+1} dy = e_\beta(0) = 1. \quad (3.4)$$

Hence, ${}^t\mathcal{X}_{\alpha,\beta}f$ is almost everywhere defined on $\mathbb{R}$, belongs to $L^1(\mathbb{R}, |x|^{2\alpha+1} dx)$ and $\|{}^t\mathcal{X}_{\alpha,\beta}f\|_{1,\alpha} \leq \|f\|_{1,\beta}$, which proves (i). Identity (3.2) follows by using (1.1), (2.1), (3.1), and Fubini's theorem. Identity (3.3) follows by applying the Dunkl transform $\mathcal{F}_\alpha$ to both its sides and by using (2.6), (3.2) and Remark 2.2. ■

Remark 3.4. From (3.2) and Remark 2.2, we deduce that the transform ${}^t\mathcal{X}_{\alpha,\beta}$ maps $L^1(\mathbb{R}, |x|^{2\beta+1} dx)$ injectively into $L^1(\mathbb{R}, |x|^{2\alpha+1} dx)$.

From [14] we have the following result.

Theorem 3.5. *The dual Dunkl–Sonine integral transform ${}^t\mathcal{X}_{\alpha,\beta}$ is an automorphism of $\mathcal{S}(\mathbb{R})$ satisfying the intertwining relation*

$${}^t\mathcal{X}_{\alpha,\beta}\Lambda_\beta f = \Lambda_\alpha {}^t\mathcal{X}_{\alpha,\beta}f, \quad f \in \mathcal{S}(\mathbb{R}).$$

Moreover ${}^t\mathcal{X}_{\alpha,\beta}$ admits the factorization

$${}^t\mathcal{X}_{\alpha,\beta}f = {}^tV_\alpha^{-1} \circ {}^tV_\beta f \quad \text{for all } f \in \mathcal{S}(\mathbb{R}),$$

where for $\gamma > -1/2$, ${}^tV_\gamma$ denotes the dual Dunkl intertwining operator given by

$${}^tV_\gamma f(y) = \frac{\Gamma(\gamma+1)}{\sqrt{\pi}\Gamma(\gamma+1/2)} \int_{|x|\geq|y|} \operatorname{sgn}(x)(x+y)(x^2-y^2)^{\gamma-1/2} f(x) dx.$$

Definition 3.6 (see [14]). The Dunkl–Sonine integral transform $\mathcal{X}_{\alpha,\beta}$ is defined for a locally bounded function f on $\mathbb{R}$ by

$$\mathcal{X}_{\alpha,\beta}f(x) = \begin{cases} \int_{-|x|}^{|x|} \mathcal{K}_{\alpha,\beta}(x,y) f(y) |y|^{2\alpha+1} dy & \text{if } x \neq 0, \\ f(0) & \text{if } x = 0. \end{cases} \quad (3.5)$$

Remark 3.7.

(i) Notice that by (3.4), $\|\mathcal{X}_{\alpha,\beta}f\|_\infty \leq \|f\|_\infty$ if $f \in L^\infty(\mathbb{R})$.

(ii) It follows from (1.1) that

$$e_\beta(\lambda x) = \mathcal{X}_{\alpha,\beta}(e_\alpha(\lambda \cdot))(x) \quad (3.6)$$

for all $\lambda \in \mathbb{C}$ and $x \in \mathbb{R}$.

Proposition 3.8.

(i) For any $f \in L^\infty(\mathbb{R})$ and $g \in L^1(\mathbb{R}, |x|^{2\beta+1}dx)$ we have the duality relation

$$\int_{\mathbb{R}} \mathcal{X}_{\alpha,\beta}f(x)g(x)|x|^{2\beta+1}dx = \int_{\mathbb{R}} f(y) {}^t\mathcal{X}_{\alpha,\beta}g(y)|y|^{2\alpha+1}dy. \quad (3.7)$$

(ii) Let $f \in L^1(\mathbb{R}, |x|^{2\beta+1}dx)$ and $g \in L^\infty(\mathbb{R})$. Then

$$\mathcal{X}_{\alpha,\beta}({}^t\mathcal{X}_{\alpha,\beta}f *_\alpha g) = f *_\beta \mathcal{X}_{\alpha,\beta}g. \quad (3.8)$$

Proof. Identity (3.7) follows by using (3.1), (3.5) and Fubini's theorem. Let us check (3.8). Let $\psi \in \mathcal{S}(\mathbb{R})$. By using (3.3), (3.7) and Fubini's theorem, we have

$$\begin{aligned} \int_{\mathbb{R}} f *_\beta \mathcal{X}_{\alpha,\beta}g(x)\psi(x)|x|^{2\beta+1}dx &= \int_{\mathbb{R}} \mathcal{X}_{\alpha,\beta}g(x)\psi *_\beta f^-(x)|x|^{2\beta+1}dx \\ &= \int_{\mathbb{R}} g(y) {}^t\mathcal{X}_{\alpha,\beta}(\psi *_\beta f^-)(y)|y|^{2\alpha+1}dy = \int_{\mathbb{R}} g(y) ({}^t\mathcal{X}_{\alpha,\beta}\psi *_\alpha {}^t\mathcal{X}_{\alpha,\beta}f^-)(y)|y|^{2\alpha+1}dy, \end{aligned}$$

where $f^-(x) = f(-x)$, $x \in \mathbb{R}$. But an easy computation shows that ${}^t\mathcal{X}_{\alpha,\beta}f^- = ({}^t\mathcal{X}_{\alpha,\beta}f)^-$. Hence,

$$\begin{aligned} \int_{\mathbb{R}} f *_\beta \mathcal{X}_{\alpha,\beta}g(x)\psi(x)|x|^{2\beta+1}dx &= \int_{\mathbb{R}} g(y) {}^t\mathcal{X}_{\alpha,\beta}\psi *_\alpha ({}^t\mathcal{X}_{\alpha,\beta}f)^-(y)|y|^{2\alpha+1}dy \\ &= \int_{\mathbb{R}} {}^t\mathcal{X}_{\alpha,\beta}f *_\alpha g(y) {}^t\mathcal{X}_{\alpha,\beta}\psi(y)|y|^{2\alpha+1}dy = \int_{\mathbb{R}} \mathcal{X}_{\alpha,\beta}({}^t\mathcal{X}_{\alpha,\beta}f *_\alpha g)(x)\psi(x)|x|^{2\beta+1}dx. \end{aligned}$$

This clearly yields the result. ■

4 Calderón's formula for the Dunkl continuous wavelet transform

Throughout this section assume $\gamma > -1/2$.

Definition 4.1. We say that a function $g \in L^2(\mathbb{R}, |x|^{2\gamma+1}dx)$ is a Dunkl wavelet of order γ , if it satisfies the admissibility condition

$$0 < C_g^\gamma := \int_0^\infty |\mathcal{F}_\gamma g(\lambda)|^2 \frac{d\lambda}{\lambda} = \int_0^\infty |\mathcal{F}_\gamma g(-\lambda)|^2 \frac{d\lambda}{\lambda} < \infty. \quad (4.1)$$

Remark 4.2.

(i) If g is real-valued we have $\mathcal{F}_\gamma g(-\lambda) = \overline{\mathcal{F}_\gamma g(\lambda)}$, so (4.1) reduces to

$$0 < C_g^\gamma := \int_0^\infty |\mathcal{F}_\gamma g(\lambda)|^2 \frac{d\lambda}{\lambda} < \infty.$$

(ii) If $0 \neq g \in L^2(\mathbb{R}, |x|^{2\gamma+1}dx)$ is real-valued and satisfies

$$\exists \eta > 0 \quad \text{such that} \quad \mathcal{F}_\gamma g(\lambda) - \mathcal{F}_\gamma g(0) = \mathcal{O}(\lambda^\eta) \quad \text{as} \quad \lambda \rightarrow 0^+$$

then (4.1) is equivalent to $\mathcal{F}_\gamma g(0) = 0$.

Note 4.3. For a function g in $L^2(\mathbb{R}, |x|^{2\gamma+1}dx)$ and for $(a, b) \in (0, \infty) \times \mathbb{R}$ we write

$$g_{a,b}^\gamma(x) := \frac{1}{a^{2\gamma+2}} T_\gamma^{-b} g_a(x),$$

where T_γ^{-b} are the generalized translation operators given by (2.3), and $g_a(x) := g(x/a)$, $x \in \mathbb{R}$.

Remark 4.4. Let $g \in L^2(\mathbb{R}, |x|^{2\gamma+1}dx)$ and $a > 0$. Then it is easily checked that $g_a \in L^2(\mathbb{R}, |x|^{2\gamma+1}dx)$, $\|g_a\|_{2,\gamma} = a^{\gamma+1} \|g\|_{2,\gamma}$, and $\mathcal{F}_\gamma(g_a)(\lambda) = a^{2\gamma+2} \mathcal{F}_\gamma(g)(a\lambda)$.

Definition 4.5. Let $g \in L^2(\mathbb{R}, |x|^{2\gamma+1}dx)$ be a Dunkl wavelet of order γ . We define for regular functions f on $\mathbb{R}$, the Dunkl continuous wavelet transform by

$$\Phi_g^\gamma(f)(a, b) := \int_{\mathbb{R}} f(x) \overline{g_{a,b}^\gamma(x)} |x|^{2\gamma+1} dx$$

which can also be written in the form

$$\Phi_g^\gamma(f)(a, b) = \frac{1}{a^{2\gamma+2}} f *_\gamma \tilde{g}_a(b),$$

where $*_\gamma$ is the generalized convolution product given by (2.4), and $\tilde{g}_a(x) := \overline{g(-x/a)}$, $x \in \mathbb{R}$.

The Dunkl continuous wavelet transform has been investigated in depth in [5] in which precise definitions, examples, and a more complete discussion of its properties can be found. We look here for a Calderón formula for this transform. We start with some technical lemmas.

Lemma 4.6. For all $f, g \in L^2(\mathbb{R}, |x|^{2\gamma+1}dx)$ and all $\psi \in \mathcal{S}(\mathbb{R})$ we have the identity

$$\int_{\mathbb{R}} f *_\gamma g(x) \mathcal{F}_\gamma^{-1} \psi(x) |x|^{2\gamma+1} dx = m_\gamma \int_{\mathbb{R}} \mathcal{F}_\gamma f(\lambda) \mathcal{F}_\gamma g(\lambda) \psi^-(\lambda) |\lambda|^{2\gamma+1} d\lambda,$$

where m_γ is given by (2.2).

Proof. Fix $g \in L^2(\mathbb{R}, |x|^{2\gamma+1}dx)$ and $\psi \in \mathcal{S}(\mathbb{R})$. For $f \in L^2(\mathbb{R}, |x|^{2\gamma+1}dx)$ put

$$S_1(f) := \int_{\mathbb{R}} f *_\gamma g(x) \mathcal{F}_\gamma^{-1} \psi(x) |x|^{2\gamma+1} dx$$

and

$$S_2(f) := m_\gamma \int_{\mathbb{R}} \mathcal{F}_\gamma f(\lambda) \mathcal{F}_\gamma g(\lambda) \psi^-(\lambda) |\lambda|^{2\gamma+1} d\lambda.$$

By (2.5), (2.6) and Theorem 2.3, we see that $S_1(f) = S_2(f)$ for each $f \in L^1 \cap L^2(\mathbb{R}, |x|^{2\gamma+1}dx)$. Moreover, by using (2.5), Hölder's inequality and Theorem 2.3 we have

$$|S_1(f)| \leq \|f *_\gamma g\|_\infty \|\mathcal{F}_\gamma^{-1} \psi\|_{1,\gamma} \leq 4 \|f\|_{2,\gamma} \|g\|_{2,\gamma} \|\mathcal{F}_\gamma^{-1} \psi\|_{1,\gamma}$$

and

$$|S_2(f)| \leq m_\gamma \|\mathcal{F}_\gamma f \mathcal{F}_\gamma g\|_{1,\gamma} \|\psi\|_\infty \leq m_\gamma \|\mathcal{F}_\gamma f\|_{2,\gamma} \|\mathcal{F}_\gamma g\|_{2,\gamma} \|\psi\|_\infty = \|f\|_{2,\gamma} \|g\|_{2,\gamma} \|\psi\|_\infty,$$

which shows that the linear functionals S_1 and S_2 are bounded on $L^2(\mathbb{R}, |x|^{2\gamma+1}dx)$. Therefore $S_1 \equiv S_2$, and the lemma is proved. $\blacksquare$

Lemma 4.7. *Let $f_1, f_2 \in L^2(\mathbb{R}, |x|^{2\gamma+1}dx)$. Then $f_1 *_\gamma f_2 \in L^2(\mathbb{R}, |x|^{2\gamma+1}dx)$ if and only if $\mathcal{F}_\gamma f_1 \mathcal{F}_\gamma f_2 \in L^2(\mathbb{R}, |x|^{2\gamma+1}dx)$ and we have*

$$\mathcal{F}_\gamma(f_1 *_\gamma f_2) = \mathcal{F}_\gamma f_1 \mathcal{F}_\gamma f_2$$

in the L^2 -case.

Proof. Suppose $f_1 *_\gamma f_2 \in L^2(\mathbb{R}, |x|^{2\gamma+1}dx)$. By Lemma 4.6 and Theorem 2.3, we have for any $\psi \in \mathcal{S}(\mathbb{R})$,

$$\begin{aligned} m_\gamma \int_{\mathbb{R}} \mathcal{F}_\gamma f_1(\lambda) \mathcal{F}_\gamma f_2(\lambda) \psi(\lambda) |\lambda|^{2\gamma+1} d\lambda &= \int_{\mathbb{R}} f_1 *_\gamma f_2(x) \mathcal{F}_\gamma^{-1} \psi^-(x) |x|^{2\gamma+1} dx \\ &= \int_{\mathbb{R}} f_1 *_\gamma f_2(x) \overline{\mathcal{F}_\gamma^{-1} \psi(x)} |x|^{2\gamma+1} dx = m_\gamma \int_{\mathbb{R}} \mathcal{F}_\gamma(f_1 *_\gamma f_2)(\lambda) \psi(\lambda) |\lambda|^{2\gamma+1} d\lambda, \end{aligned}$$

which shows that $\mathcal{F}_\gamma f_1 \mathcal{F}_\gamma f_2 = \mathcal{F}_\gamma(f_1 *_\gamma f_2)$. Conversely, if $\mathcal{F}_\gamma f_1 \mathcal{F}_\gamma f_2 \in L^2(\mathbb{R}, |x|^{2\gamma+1}dx)$, then by Lemma 4.6 and Theorem 2.3, we have for any $\psi \in \mathcal{S}(\mathbb{R})$,

$$\begin{aligned} \int_{\mathbb{R}} f_1 *_\gamma f_2(x) \mathcal{F}_\gamma^{-1} \psi(x) |x|^{2\gamma+1} dx &= m_\gamma \int_{\mathbb{R}} \mathcal{F}_\gamma f_1(\lambda) \mathcal{F}_\gamma f_2(\lambda) \overline{\psi(\lambda)} |\lambda|^{2\gamma+1} d\lambda \\ &= \int_{\mathbb{R}} \mathcal{F}_\gamma^{-1}(\mathcal{F}_\gamma f_1 \mathcal{F}_\gamma f_2)(x) \mathcal{F}_\gamma^{-1} \psi(x) |x|^{2\gamma+1} dx, \end{aligned}$$

which shows, in view of Theorem 2.4, that $f_1 *_\gamma f_2 = \mathcal{F}_\gamma^{-1}(\mathcal{F}_\gamma f_1 \mathcal{F}_\gamma f_2)$. This achieves the proof of Lemma 4.7. $\blacksquare$

A combination of Lemma 4.7 and Theorem 2.3 gives us the following.

Lemma 4.8. *Let $f_1, f_2 \in L^2(\mathbb{R}, |x|^{2\gamma+1}dx)$. Then*

$$\int_{\mathbb{R}} |f_1 *_\gamma f_2(x)|^2 |x|^{2\gamma+1} dx = m_\gamma \int_{\mathbb{R}} |\mathcal{F}_\gamma f_1(\lambda)|^2 |\mathcal{F}_\gamma f_2(\lambda)|^2 |\lambda|^{2\gamma+1} d\lambda,$$

where both sides are finite or infinite.

Lemma 4.9. *Let $g \in L^2(\mathbb{R}, |x|^{2\gamma+1} dx)$ be a Dunkl wavelet of order γ such that $\mathcal{F}_\gamma g \in L^\infty(\mathbb{R})$. For $0 < \varepsilon < \delta < \infty$ define*

$$G_{\varepsilon, \delta}(x) := \frac{1}{C_g^\gamma} \int_\varepsilon^\delta g_a *_\gamma \tilde{g}_a(x) \frac{da}{a^{4\gamma+5}} \quad (4.2)$$

and

$$K_{\varepsilon, \delta}(\lambda) := \frac{1}{C_g^\gamma} \int_\varepsilon^\delta |\mathcal{F}_\gamma g(a\lambda)|^2 \frac{da}{a}. \quad (4.3)$$

Then

$$G_{\varepsilon, \delta} \in L^2(\mathbb{R}, |x|^{2\gamma+1} dx), \quad K_{\varepsilon, \delta} \in (L^1 \cap L^2)(\mathbb{R}, |x|^{2\gamma+1} dx), \quad (4.4)$$

and

$$\mathcal{F}_\gamma(G_{\varepsilon, \delta}) = K_{\varepsilon, \delta}.$$

Proof. Using Schwarz inequality for the measure $\frac{da}{a^{4\gamma+5}}$ we obtain

$$|G_{\varepsilon, \delta}(x)|^2 \leq \frac{1}{(C_g^\gamma)^2} \left(\int_\varepsilon^\delta \frac{da}{a^{4\gamma+5}} \right) \int_\varepsilon^\delta |g_a *_\gamma \tilde{g}_a(x)|^2 \frac{da}{a^{4\gamma+5}},$$

so

$$\int_{\mathbb{R}} |G_{\varepsilon, \delta}(x)|^2 |x|^{2\gamma+1} dx \leq \frac{1}{(C_g^\gamma)^2} \left(\int_\varepsilon^\delta \frac{da}{a^{4\gamma+5}} \right) \int_\varepsilon^\delta \int_{\mathbb{R}} |g_a *_\gamma \tilde{g}_a(x)|^2 |x|^{2\gamma+1} dx \frac{da}{a^{4\gamma+5}}.$$

By Theorem 2.3, Lemma 4.8, and Remark 4.4, we have

$$\begin{aligned} \int_{\mathbb{R}} |g_a *_\gamma \tilde{g}_a(x)|^2 |x|^{2\gamma+1} dx &= m_\gamma \int_{\mathbb{R}} |\mathcal{F}_\gamma(g_a)(\lambda)|^4 |\lambda|^{2\gamma+1} d\lambda \\ &\leq m_\gamma \|\mathcal{F}_\gamma(g_a)\|_\infty^2 \int_{\mathbb{R}} |\mathcal{F}_\gamma(g_a)(\lambda)|^2 |\lambda|^{2\gamma+1} d\lambda \\ &= \|\mathcal{F}_\gamma(g_a)\|_\infty^2 \|g_a\|_{2, \gamma}^2 = a^{6\gamma+6} \|\mathcal{F}_\gamma g\|_\infty^2 \|g\|_{2, \gamma}^2. \end{aligned}$$

Hence

$$\int_{\mathbb{R}} |G_{\varepsilon, \delta}(x)|^2 |x|^{2\gamma+1} dx \leq \frac{\|\mathcal{F}_\gamma g\|_\infty^2 \|g\|_{2, \gamma}^2}{(C_g^\gamma)^2} \left(\int_\varepsilon^\delta a^{2\gamma+1} da \right) \left(\int_\varepsilon^\delta \frac{da}{a^{4\gamma+5}} \right) < \infty.$$

The second assertion in (4.4) is easily checked. Let us calculate $\mathcal{F}_\gamma(G_{\varepsilon, \delta})$. Fix $x \in \mathbb{R}$. From Theorem 2.3 and Lemma 4.7 we get

$$g_a *_\gamma \tilde{g}_a(x) = m_\gamma \int_{\mathbb{R}} |\mathcal{F}_\gamma(g_a)(\lambda)|^2 e_\gamma(i\lambda x) |\lambda|^{2\gamma+1} d\lambda,$$

so

$$G_{\varepsilon, \delta}(x) = \frac{m_\gamma}{C_g^\gamma} \int_\varepsilon^\delta \left(\int_{\mathbb{R}} |\mathcal{F}_\gamma(g_a)(\lambda)|^2 e_\gamma(i\lambda x) |\lambda|^{2\gamma+1} d\lambda \right) \frac{da}{a^{4\gamma+5}}.$$

As $|e_\gamma(iz)| \leq 1$ for all $z \in \mathbb{R}$ (see [12]), we deduce by Theorem 2.3 that

$$m_\gamma \int_\varepsilon^\delta \int_{\mathbb{R}} |\mathcal{F}_\gamma(g_a)(\lambda)|^2 |e_\gamma(i\lambda x)| |\lambda|^{2\gamma+1} d\lambda \frac{da}{a^{4\gamma+5}}$$

$$\leq \int_{\varepsilon}^{\delta} \|g_a\|_{2,\gamma}^2 \frac{da}{a^{4\gamma+5}} = \|g\|_{2,\gamma}^2 \int_{\varepsilon}^{\delta} \frac{da}{a^{2\gamma+3}} < \infty.$$

Hence, applying Fubini's theorem, we find that

$$\begin{aligned} G_{\varepsilon,\delta}(x) &= m_{\gamma} \int_{\mathbb{R}} \left(\frac{1}{C_g^{\gamma}} \int_{\varepsilon}^{\delta} |\mathcal{F}_{\gamma} g(a\lambda)|^2 \frac{da}{a} \right) e_{\gamma}(i\lambda x) |\lambda|^{2\gamma+1} d\lambda \\ &= m_{\gamma} \int_{\mathbb{R}} K_{\varepsilon,\delta}(\lambda) e_{\gamma}(i\lambda x) |\lambda|^{2\gamma+1} d\lambda \end{aligned}$$

which completes the proof. ■

We can now state the main result of this section.

Theorem 4.10 (Calderón's formula). *Let $g \in L^2(\mathbb{R}, |x|^{2\gamma+1} dx)$ be a Dunkl wavelet of order γ such that $\mathcal{F}_{\gamma} g \in L^{\infty}(\mathbb{R})$. Then for $f \in L^2(\mathbb{R}, |x|^{2\gamma+1} dx)$ and $0 < \varepsilon < \delta < \infty$, the function*

$$f^{\varepsilon,\delta}(x) := \frac{1}{C_g^{\gamma}} \int_{\varepsilon}^{\delta} \int_{\mathbb{R}} \Phi_g^{\gamma}(f)(a, b) g_{a,b}(x) |b|^{2\gamma+1} db \frac{da}{a}$$

belongs to $L^2(\mathbb{R}, |x|^{2\gamma+1} dx)$ and satisfies

$$\lim_{\varepsilon \rightarrow 0, \delta \rightarrow \infty} \|f^{\varepsilon,\delta} - f\|_{2,\gamma} = 0. \quad (4.5)$$

Proof. It is easily seen that

$$f^{\varepsilon,\delta} = f *_{\gamma} G_{\varepsilon,\delta},$$

where $G_{\varepsilon,\delta}$ is given by (4.2). It follows by Lemmas 4.7 and 4.9 that $f^{\varepsilon,\delta} \in L^2(\mathbb{R}, |x|^{2\gamma+1} dx)$ and $\mathcal{F}_{\gamma}(f^{\varepsilon,\delta}) = \mathcal{F}_{\gamma}(f) K_{\varepsilon,\delta}$, where $K_{\varepsilon,\delta}$ is as in (4.3). From this and Theorem 2.3 we obtain

$$\begin{aligned} \|f^{\varepsilon,\delta} - f\|_{2,\gamma}^2 &= m_{\gamma} \int_{\mathbb{R}} |\mathcal{F}_{\gamma}(f^{\varepsilon,\delta} - f)(\lambda)|^2 |\lambda|^{2\gamma+1} d\lambda \\ &= m_{\gamma} \int_{\mathbb{R}} |\mathcal{F}_{\gamma} f(\lambda)|^2 (1 - K_{\varepsilon,\delta}(\lambda))^2 |\lambda|^{2\gamma+1} d\lambda. \end{aligned}$$

But by (4.1) we have

$$\lim_{\varepsilon \rightarrow 0, \delta \rightarrow \infty} K_{\varepsilon,\delta}(\lambda) = 1, \quad \text{for almost all } \lambda \in \mathbb{R}.$$

So (4.5) follows from the dominated convergence theorem. ■

Another pointwise inversion formula for the Dunkl wavelet transform, proved in [5], is as follows.

Theorem 4.11. *Let $g \in L^2(\mathbb{R}, |x|^{2\gamma+1} dx)$ be a Dunkl wavelet of order γ . If both f and $\mathcal{F}_{\gamma} f$ are in $L^1(\mathbb{R}, |x|^{2\gamma+1} dx)$ then we have*

$$f(x) = \frac{1}{C_g^{\gamma}} \int_0^{\infty} \left(\int_{\mathbb{R}} \Phi_g^{\gamma}(f)(a, b) g_{a,b}(x) |b|^{2\gamma+1} db \right) \frac{da}{a}, \quad a.e.,$$

where, for each $x \in \mathbb{R}$, both the inner integral and the outer integral are absolutely convergent, but possibly not the double integral.

5 Inversion of the dual Dunkl–Sonine transform using Dunkl wavelets

From now on assume $\beta > \alpha > -1/2$. In order to invert the dual Dunkl–Sonine transform, we need the following two technical lemmas.

Lemma 5.1. *Let $0 \neq g \in L^1 \cap L^2(\mathbb{R}, |x|^{2\alpha+1}dx)$ such that $\mathcal{F}_\alpha g \in L^1(\mathbb{R}, |x|^{2\alpha+1}dx)$ and satisfying*

$$\exists \eta > \beta - 2\alpha - 1 \quad \text{such that} \quad \mathcal{F}_\alpha g(\lambda) = \mathcal{O}(|\lambda|^\eta) \quad \text{as} \quad \lambda \rightarrow 0. \quad (5.1)$$

Then $\mathcal{X}_{\alpha,\beta}g \in L^2(\mathbb{R}, |x|^{2\beta+1}dx)$ and

$$\mathcal{F}_\beta(\mathcal{X}_{\alpha,\beta}g)(\lambda) = \frac{m_\alpha}{m_\beta} \frac{\mathcal{F}_\alpha g(\lambda)}{|\lambda|^{2(\beta-\alpha)}}.$$

Proof. By Theorem 2.3 we have

$$g(x) = m_\alpha \int_{\mathbb{R}} \mathcal{F}_\alpha g(\lambda) e_\alpha(i\lambda x) |\lambda|^{2\alpha+1} d\lambda, \quad \text{a.e.}$$

So using (3.6), we find that

$$\mathcal{X}_{\alpha,\beta}g(x) = m_\beta \int_{\mathbb{R}} h_{\alpha,\beta}(\lambda) e_\beta(i\lambda x) |\lambda|^{2\beta+1} d\lambda, \quad \text{a.e.} \quad (5.2)$$

with

$$h_{\alpha,\beta}(\lambda) := \frac{m_\alpha}{m_\beta} \frac{\mathcal{F}_\alpha g(\lambda)}{|\lambda|^{2(\beta-\alpha)}}.$$

Clearly, $h_{\alpha,\beta} \in L^1(\mathbb{R}, |x|^{2\beta+1}dx)$. So it suffices, in view of (5.2) and Theorem 2.3, to prove that $h_{\alpha,\beta}$ belongs to $L^2(\mathbb{R}, |x|^{2\beta+1}dx)$. We have

$$\begin{aligned} \int_{\mathbb{R}} |h_{\alpha,\beta}(\lambda)|^2 |\lambda|^{2\beta+1} d\lambda &= \left(\frac{m_\alpha}{m_\beta} \right)^2 \int_{\mathbb{R}} |\lambda|^{4\alpha-2\beta+1} |\mathcal{F}_\alpha g(\lambda)|^2 d\lambda \\ &= \left(\frac{m_\alpha}{m_\beta} \right)^2 \left(\int_{|\lambda| \leq 1} + \int_{|\lambda| \geq 1} \right) |\lambda|^{4\alpha-2\beta+1} |\mathcal{F}_\alpha g(\lambda)|^2 d\lambda := I_1 + I_2. \end{aligned}$$

By (5.1) there is a positive constant k such that

$$I_1 \leq k \int_{|\lambda| \leq 1} |\lambda|^{2\eta+4\alpha-2\beta+1} d\lambda = \frac{k}{\eta + 2\alpha - \beta + 1} < \infty.$$

From Theorem 2.3, it follows that

$$\begin{aligned} I_2 &= \left(\frac{m_\alpha}{m_\beta} \right)^2 \int_{|\lambda| \geq 1} |\lambda|^{2(\alpha-\beta)} |\mathcal{F}_\alpha g(\lambda)|^2 |\lambda|^{2\alpha+1} d\lambda \\ &\leq \left(\frac{m_\alpha}{m_\beta} \right)^2 \int_{|\lambda| \geq 1} |\mathcal{F}_\alpha g(\lambda)|^2 |\lambda|^{2\alpha+1} d\lambda \leq \left(\frac{m_\alpha}{m_\beta} \right)^2 \|\mathcal{F}_\alpha g\|_{2,\alpha}^2 = \frac{m_\alpha}{(m_\beta)^2} \|g\|_{2,\alpha}^2 < \infty \end{aligned}$$

which ends the proof. ■

Lemma 5.2. *Let $0 \neq g \in L^1 \cap L^2(\mathbb{R}, |x|^{2\alpha+1}dx)$ be real-valued such that $\mathcal{F}_\alpha g \in L^1(\mathbb{R}, |x|^{2\alpha+1}dx)$ and satisfying*

$$\exists \eta > 2(\beta - \alpha) \quad \text{such that} \quad \mathcal{F}_\alpha g(\lambda) = \mathcal{O}(\lambda^\eta) \quad \text{as} \quad \lambda \rightarrow 0^+. \quad (5.3)$$

Then $\mathcal{X}_{\alpha,\beta}g \in L^2(\mathbb{R}, |x|^{2\beta+1}dx)$ is a Dunkl wavelet of order β and $\mathcal{F}_\beta(\mathcal{X}_{\alpha,\beta}g) \in L^\infty(\mathbb{R})$.

Proof. By combining (5.3) and Lemma 5.1 we see that $\mathcal{X}_{\alpha,\beta}g \in L^2(\mathbb{R}, |x|^{2\beta+1}dx)$, $\mathcal{F}_\beta(\mathcal{X}_{\alpha,\beta}g)$ is bounded and

$$\mathcal{F}_\beta(\mathcal{X}_{\alpha,\beta}g)(\lambda) = \mathcal{O}(\lambda^{\eta-2(\beta-\alpha)}) \quad \text{as} \quad \lambda \rightarrow 0^+.$$

Thus, in view of Remark 4.2, $\mathcal{X}_{\alpha,\beta}g$ satisfies the admissibility condition (4.1) for $\gamma = \beta$. ■

Remark 5.3. In view of Remark 4.2, each function satisfying the conditions of Lemma 5.1 is a Dunkl wavelet of order α .

Lemma 5.4. *Let g be as in Lemma 5.2. Then for all $f \in L^1(\mathbb{R}, |x|^{2\beta+1}dx)$ we have*

$$\Phi_{\mathcal{X}_{\alpha,\beta}g}^\beta(f)(a, b) = \frac{1}{a^{2(\beta-\alpha)}} \mathcal{X}_{\alpha,\beta} [\Phi_g^\alpha({}^t\mathcal{X}_{\alpha,\beta}f)(a, \cdot)](b).$$

Proof. By Definition 4.5 we have

$$\Phi_{\mathcal{X}_{\alpha,\beta}g}^\beta(f)(a, b) = \frac{1}{a^{2\beta+2}} f *_\beta (\widetilde{\mathcal{X}_{\alpha,\beta}g})_a(b).$$

But $(\widetilde{\mathcal{X}_{\alpha,\beta}g})_a = \mathcal{X}_{\alpha,\beta}(\widetilde{g}_a)$ by virtue of (1.2) and (3.5). So using (3.8) we find that

$$\begin{aligned} \Phi_{\mathcal{X}_{\alpha,\beta}g}^\beta(f)(a, b) &= \frac{1}{a^{2\beta+2}} f *_\beta [\mathcal{X}_{\alpha,\beta}(\widetilde{g}_a)](b) \\ &= \frac{1}{a^{2\beta+2}} \mathcal{X}_{\alpha,\beta} [{}^t\mathcal{X}_{\alpha,\beta}f *_\alpha \widetilde{g}_a](b) = \frac{1}{a^{2(\beta-\alpha)}} \mathcal{X}_{\alpha,\beta} [\Phi_g^\alpha({}^t\mathcal{X}_{\alpha,\beta}f)(a, \cdot)](b), \end{aligned}$$

which gives the desired result. ■

Combining Theorems 4.10, 4.11 with Lemmas 5.2, 5.4 we get

Theorem 5.5. *Let g be as in Lemma 5.2. Then we have the following inversion formulas for the dual Dunkl–Sonine transform:*

(i) *If both f and $\mathcal{F}_\beta f$ are in $L^1(\mathbb{R}, |x|^{2\beta+1}dx)$ then for almost all $x \in \mathbb{R}$ we have*

$$f(x) = \frac{1}{C_{\mathcal{X}_{\alpha,\beta}g}^\beta} \int_0^\infty \left(\int_{\mathbb{R}} \mathcal{X}_{\alpha,\beta} [\Phi_g^\alpha({}^t\mathcal{X}_{\alpha,\beta}f)(a, \cdot)](b) (\mathcal{X}_{\alpha,\beta}g)_{a,b}^\beta(x) |b|^{2\beta+1} db \right) \frac{da}{a^{2(\beta-\alpha)+1}}.$$

(ii) *For $f \in L^1 \cap L^2(\mathbb{R}, |x|^{2\beta+1}dx)$ and $0 < \varepsilon < \delta < \infty$, the function*

$$f^{\varepsilon,\delta}(x) := \frac{1}{C_{\mathcal{X}_{\alpha,\beta}g}^\beta} \int_\varepsilon^\delta \int_{\mathbb{R}} \mathcal{X}_{\alpha,\beta} [\Phi_g^\alpha({}^t\mathcal{X}_{\alpha,\beta}f)(a, \cdot)](b) (\mathcal{X}_{\alpha,\beta}g)_{a,b}^\beta(x) |b|^{2\beta+1} db \frac{da}{a^{2(\beta-\alpha)+1}}$$

satisfies

$$\lim_{\varepsilon \rightarrow 0, \delta \rightarrow \infty} \|f^{\varepsilon,\delta} - f\|_{2,\beta} = 0.$$

Acknowledgements

The author is grateful to the referees and editors for careful reading and useful comments.

References

- [1] de Jeu M.F.E., The Dunkl transform, *Invent. Math.* **113** (1993), 147–162.
- [2] Dunkl C.F., Differential-difference operators associated with reflections groups, *Trans. Amer. Math. Soc.* **311** (1989), 167–183.
- [3] Dunkl C.F., Integral kernels with reflection group invariance, *Canad. J. Math.* **43** (1991), 1213–1227.
- [4] Dunkl C.F., Hankel transforms associated to finite reflection groups, *Contemp. Math.* **138** (1992), 123–138.
- [5] Jouini A., Dunkl wavelets and applications to inversion of the Dunkl intertwining operator and its dual, *Int. J. Math. Math. Sci.* **6** (2004), 285–293.
- [6] Kamefuchi S., Ohnuki Y., Quantum field theory and parastatistics, Springer-Verlag, Berlin, 1982.
- [7] Mourou M.A., Trimèche K., Calderón’s formula associated with a differential operator on $(0, \infty)$ and inversion of the generalized Abel transform, *J. Fourier Anal. Appl.* **4** (1998), 229–245.
- [8] Mourou M.A., Trimèche K., Inversion of the Weyl integral transform and the Radon transform on $\mathbb{R}^n$ using generalized wavelets, *Monatsh. Math.* **126** (1998), 73–83.
- [9] Mourou M.A., Trimèche K., Calderon’s reproducing formula related to the Dunkl operator on the real line, *Monatsh. Math.* **136** (2002), 47–65.
- [10] Mourou M.A., Trimèche K., Transmutation operators and Paley–Wiener associated with a singular differential-difference operator on the real line, *Anal. Appl. (Singap.)* **1** (2003), 43–70.
- [11] Rosenblum M., Generalized Hermite polynomials and the Bose-like oscillator calculus, in Nonselfadjoint Operators and Related Topics (Beer Sheva, 1992), *Oper. Theory Adv. Appl.*, Vol. 73, Birkhäuser, Basel, 1994, 369–396, math.CA/9307224.
- [12] Rösler M., Bessel-type signed hypergroups on $\mathbb{R}$, in Probability Measures on Groups and Related Structures, XI (Oberwolfach, 1994), Editors H. Heyer and A. Mukherjea, Oberwolfach, 1994, World Sci. Publ., River Edge, NJ, 1995, 292–304.
- [13] Soltani F., L^p -Fourier multipliers for the Dunkl operator on the real line, *J. Funct. Anal.* **209** (2004), 16–35.
- [14] Soltani F., Sonine transform associated to the Dunkl kernel on the real line, *SIGMA* **4** (2008), 092, 14 pages, arXiv:0812.4666.
- [15] Xu Y., An integral formula for generalized Gegenbauer polynomials and Jacobi polynomials, *Adv. in Appl. Math.* **29** (2002), 328–343.
- [16] Yang L.M., A note on the quantum rule of the harmonic oscillator, *Phys. Rev.* **84** (1951), 788–790.